



AIAA-98-1042

A Survey of Spacecraft Charging Events

Shu T. Lai

Air Force Research Laboratory

29 Randolph Road

Hanscom AFB, MA 01731-3010

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

**36th Aerospace Sciences
Meeting & Exhibit**

January 12-15, 1998 / Reno, NV

19980325 085

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 17 March 1998	3. REPORT TYPE AND DATES COVERED Reprint		
4. TITLE AND SUBTITLE A Survey of Spacecraft Charging Events		5. FUNDING NUMBERS PR 1010 TA RS WU 01		
6. AUTHOR(S) Shu T. Lai				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory/VSBS 29 Randolph Road Hanscom AFB, MA 01731-3010		8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-VS-HA-TR-98-0036		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES Reprinted from 36th Aerospace Sciences Meeting & Exhibit, 12-15 January 1998, Reno, NV				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) This is a survey, or overview of spacecraft charging events. The emphasis is on concepts and physics, rather than enumerative accounts of projects or detailed statistics of spacecraft. <div style="text-align: right; transform: rotate(-15deg); font-weight: bold;">DTIC QUALITY INSPECTED 2</div>				
14. SUBJECT TERMS Spacecraft charging Spacecraft anomalies Bulk charging		15. NUMBER OF PAGES		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR	

A Survey of Spacecraft Charging Events

Shu T. Lai
Air Force Research Laboratory
Hanscom AFB., MA 01731-3010

This is a survey, or overview, of spacecraft charging events. The emphasis is on concepts and physics, rather than enumerative accounts of projects or detailed statistics of spacecraft.

1. Spacecraft Surface Charging

When an object is placed in a plasma, the object is likely charged negatively relative to the plasma. The reason is because the plasma electrons, being much lighter and faster than the ions, have a larger flux than that of ions. As a result, more electrons impact on the object surface than ions, resulting in a negative surface potential. At equilibrium, the negative potential, repelling electrons and attracting ions, achieves an equality of electron and ion currents.

2. Adverse Effects

Spacecraft surface charging may affect the scientific measurements onboard, such as measurements of the ambient plasma density and energy, the ambient electric fields, and geomagnetic fields. It may also affect contaminations such as ion deposition on the surfaces of spacecraft and mirrors. In extreme situations, it may cause stray signals in telemetry, undesirable signals in circuits, or even erroneous commands in navigation systems.

3. Where does Spacecraft Charging Occur?

Surface charging occurs not only in space but also in the laboratory. The space physicist is mostly concerned with natural charging at geosynchronous altitudes and low Earth orbits (LEO). Deep or bulk charging occurs in the Radiation Belts. Artificial charging can occur during beam emissions from a spacecraft or when the spacecraft has a long tether.

At or near the geosynchronous altitudes, spacecraft charging is important not only because the ambient plasma density is low and energy sometimes high but also there are many spacecraft in that region. At these altitudes, the ion flux is often two orders of magnitude less than that of electrons. Charging up to several kV can occur. The spacecraft SCATHA was launched in 1979 for dedicated study of spacecraft charging at these altitudes.

There is a local time effect at the geosynchronous altitudes. Due to the existence of cross-tail electric fields, the plasma coming from the magnetic tail towards the Earth would drift towards the midnight to dawn sector. Since energetic plasmas often come the tail, the midnight to dawn sector is where charging likely occurs.

At LEO altitudes, charging is at low level, Volts, only. This is because the ambient plasma in this region is dense and not energetic. If the surface potential tries to increase, the opposite charges would be attracted in abundance to prevent any high potential formation.

Since the ions in LEO are slower than a spacecraft, there is a void of ions in the spacecraft wake, where the potential tends to be negative.

The only significant natural charging region in the LEO environment is the auroral zone where the electrons are often directional and energetic.

4. Current Balance

The surface potential at equilibrium is determined by the balance of currents, according to

Kirchhoff's circuital law. Besides ambient plasma electron and ion currents, photoelectron emission plays an important role for high altitude charging in sunlight. Artificial beam emissions of electrons, ions, or both can greatly control the surface potential if the beam currents exceed the ambient currents.

Secondary and backscattered electron emission coefficients are properties of surface materials and are functions of the primary electron energy. For a given surface, if the sum of these coefficients exceeds unity, positive charging occurs. This likely occurs for many materials when the primary electron energies are in the range of about 40 to 1000 eV. The surface condition and the angle of electron impact can also affect the emissions. The coefficients play crucial roles in charging.

5. Onset of Spacecraft Charging

When the ambient plasma is quiet, it can be described approximately by a Maxwellian distribution $f(E)$.

$$f(E) = n(m/2kT)^{1/2} \exp(-E/kT)$$

The current balance equation is given by

$$\int dE E [\delta(E) + \eta(E)] f(E) = \int dE f(E)$$

where δ and η are the secondary and backscattered electron emission coefficients respectively. For Maxwellian distributions, the solution of this equation gives a critical temperature T_* [Lai, et al., 1982, 1983; Laframboise, 1982].

TABLE 1. Critical Temperatures (eV)

Material	Isotropic	Normal
Kapton	800	500
Teflon	2100	1400
Cu-Be	2100	1300
Cu-Be (Activated)	5300	3700
Silver	2700	1200
Gold	4900	2900
MgO	3600	2500
SiO ₂	2600	1700




Figure 1. ATS-5 Spacecraft charging data [Rubin et al., 1980]. The existence of a critical temperature manifests.

When the ambient plasma temperature T is below T_* , no charging occurs. As T reaches and exceeds T_* , onset of charging occurs. The charging events on ATS-5 agree with the theoretical predictions [Lai, 1991a].

6. Charging in a Double Maxwellian Plasma

At geosynchronous altitudes, there are occasionally energetic (multi-keV) plasma clouds arriving from the Sun or the geomagnetic tail. When they arrive, the local plasma distribution becomes a double Maxwellian approximately:

$$f(E) = n_1(m/kT_1)^{1/2} \exp(-E/kT_1) + n_2(m/kT_2)^{1/2} \exp(-E/kT_2)$$

The first one is often of low energy (up to 1 keV approximately) while the second one many keVs. Since secondary and backscattered emissions are important in the energy range up to about 1 keV for primary electrons, the first Maxwellian distribution often favors positive charging. At higher primary electron energies, the emission fluxes are small. Thus, the second Maxwellian distribution favors negative charging. Whether spacecraft charging occurs depends on the competition between the two Maxwellians. On Day 114 of SCATHA, as the first Maxwellian diminishes steadily, the second one wins

eventually, with onset of charging. This event happens to be a triple-root jump in spacecraft potential [Lai, 1991a, 1991b].




Fig.2 Double Maxwellian parameters on Day 114. [Lai, 1991a]




Fig.3 Triple-root domain on Day 114. [Lai, 1991a]

7. Differential Charging

Since different surface materials have different properties such as secondary and backscattered emissions, a spacecraft covered with different pieces of surface materials may suffer from differential charging. It may occur naturally or during beam emissions. When electron beams are emitted, electrons are drawn from the spacecraft ground

connected to the beam device. As a result, the dielectric surfaces and the ground become differentially charged.

Differential charging is undesirable because it tends to induce interactions between surfaces of different potentials. Electrons may flow from one surface to another, disturbing scientific measurements onboard or causing anomalies to electronic circuits. In extreme cases, sudden developments of differential charging may lead to sudden discharges, generating harmful electromagnetic pulses.

The amount of charge Q stored between the surface of a thin dielectric layer and the underlying spacecraft ground, for example, is given by:

$$Q = (eA/d)\Delta\phi$$

where d is the thickness, A surface area, $\Delta\phi$ the potential difference. The amount Q may be large for a thin layer, such as a coat of paint.

8. Damage to Onboard Electronics

A most dramatic event occurred on SCATHA. It was the damage of the instruments S2-1 and S2-2 during the emission of a 13 mA, 1.5 keV, electron beam. The spacecraft was charged to near beam energy. The instruments were oppositely deployed booms of about 3m long for measuring the sheath potential. The S2-1 was destroyed when one-third of its boom was in shadow and two-thirds in sunlight. Half rotation period later, S2-2 rotated into the same shadow and was destroyed similarly. This event has been fully documented [Cohen et al., 1981] but the exact cause was never found.

9. Anomalies due to Surface Charging

Anomalies occur occasionally during surface charging. For a recent example, a lockup of a microprocessor unit on a DMSP satellite [Anderson and Koons, 1996] occurred during a surface charging event of about 500 V in the auroral zone. The layered structure of the thermal dielectric blankets on that spacecraft had a large capacitance, and, a discharge in the thermal blankets probably caused

the anomaly [Anderson and Koons, 1996].

Vampola [1987] has presented a survey discussion on many anomaly events, especially those due to deep dielectric charging.

10. Deep Dielectric Charging

The Sun controls the Earth's space weather. Occasionally the Sun emits an energetic plasma cloud called a solar coronal mass ejection. The plasma in the cloud often reach MeV and beyond. Since energetic electrons and ions can penetrate deep into materials, they deposit inside. The depth of deposition depends on the energy of the incoming electron or ion. At MeV energy, electrons penetrate deeper than ions. A MeV plasma impacting on a spacecraft surface material would form a double layer, the deeper layer being that of electrons. For dielectrics, the deposited charges can stay for hours and even days. They migrate and escape slowly because of the low conductivity of dielectrics, and may cause little harm if the flux is low or the neutralization process is slow.

11. Effects of Deep Dielectric Charging

Deep dielectric charging gives typically very low spacecraft surface potential. This is mainly because the energetic (MeV) fluxes are small. Furthermore, the electrons, although more abundant, are deposited deeper than the ions, which are deposited nearer the surface.

Violet and Frederickson [1993] reported that many anomaly events occurred on CRRES during low spacecraft surface potentials. They occurred when CRRES was in the Radiation Belts where deep dielectric charging can take place.

Although the MeV fluxes are usually low, the density of charges deposited can accumulate over time. Local electric fields can reach up to 10^6 to 10^8 V/m [Hastings and Garrett, 1986] which may cause breakdowns.

Despite their low fluxes, some MeV electrons may penetrate into the electronics onboard. It is known that there is significant correlation between high fluences of electron depositions and anomalies in some types of electronic devices.




Fig.4 Deep dielectric charging on CRRES. [Violet and Frederickson, 1992].




Fig.5 Deep dielectric charging on DRAδ. [Wrenn, 1995]

At energies of about 100 MeVs or higher, ions can penetrate into materials deeper than electrons and produce cascades of ionization. Such energetic particle fluxes are very small.



For more on spacecraft anomalies due to the radiation environment, see, for example, Lauriente et al. [1998].

12. AT&T Telstar 401

On Jan 11, 1997, AT&T Telstar 401 at geosynchronous altitudes failed (EOS, 49-51, 1997; www-istp.gsfc.nasa.gov/istp/cloud_jan97/att.html and umbra.nascom.nasa.gov/istp/SHINE_report.html). A solar coronal mass ejection cloud of energetic plasma (MeV) arrived at the geosynchronous region

a day earlier and the cloud passage lasted a day. The Telstar failure occurred after the passage of the cloud. While the exact mechanisms of the failure are still at large, a common view seems to be deep dielectric charging.

The following figures show the electron and ion spectra in a geosynchronous vicinity of Telstar 401 on Jan 10, 1977. The depths of deposition are calculated by using the typical energy-deposition relation in Hastings and Garrett [1996].



13. Discussions

The exact mechanism of the failure may never be found. A few comments are as follows:

(1) The spatial distribution of electron and ion depositions is time dependent. Since the electron flux is orders of magnitude higher than that of ions, the electron layer is formed first. After a cloud passage, the ambient plasma is not energetic but more dense. The ion layer is then formed near the surface. Thus, a discharge is more likely after a storm passage.

(2) Electrons stay inside dielectrics for days. It takes time for them to migrate. However, what triggers them to avalanche during the migration is still unknown.

A case in point is Vampola's [1987] citation of Robbins [1979], who reported that anomalies on Meteosat occurred days after the passage of storms, not during them.

Wrenn [1995; 1996] show good correlation between high fluences and anomalies on DRAδ. However, the figures are not plotted in fine time scales.

The Telstar 401 event reminds us of the failure of another geosynchronous satellite, ANIK E1, which occurred after two weeks of energetic conditions [Baker, et al., 1997].

Acknowledgment

The author wish to thank Gary Mullen for initiating my interest in the 1997 January 10-11 storm and Geoff Reeves for the storm data.

References

- Anderson P.C. and H.C. Koons, Spacecraft charging anomaly on a low-altitude satellite in an aurora, J. Spacecraft & Rockets, Vol. 33, No.5, 734-738. 1996.
- Baker, D.N., et al., An assessment of space environmental conditions during the recent Anik E1 spacecraft operational failure, ISTP CDHF Newsletter Vol.6-2, 1997.

Cohen H.A., et al, P78-2 satellite and payload

responses to electron beam operations on March 30, 1979, *Proc. Spacecraft Charging Technology Conf.*, C.P. Pike, N.J. Stevens, eds, NASA-CP-2181, pp.509-559, 1981.

Frederickson, A.R., E.G. Holeman, and E.G. Mullen, Characteristics of spontaneous electrical discharging of various insulators in space radiations, *IEEE Trans. Nucl. Sci.*, 39, 1773-1782, 1992.

Hastings, D. and H. Garrett, *Spacecraft-Environment Interactions*, Cambridge University Press, 1996.

Garrett, H.B., The charging of spacecraft surfaces, *Rev. Geophys. Space Phys.*, Vol.19, 577-616, 1981.

Lai, S.T., M.S. Gussenhoven and H.A. Cohen, Range of electron energy spectrum responsible for spacecraft charging, *AGU Spring Mtg*, Philadelphia, PA, May, 1982; *EOS*, Vol.63, No.18, p.421, 1982.

Lai, S.T., M.S. Gussenhoven, H.A. Cohen, The concepts of critical temperature and energy cutoff of ambient electrons in high-voltage charging of spacecrafts, in *Spacecraft / Plasma Interactions and Their Influence on Field and Particle Measurements*, A. Pedersen, D. Guyenne, J. Hunt, eds., ESA-SP-198, 169-175, 1983. (DTIC ADA137746)

Lai, S.T., An overview of electron and ion beam effects in charging and discharging of spacecraft, *IEEE Trans. Nuclear Sci.*, 36, 2027-2032, 1989.

Lai, S.T., Theory and observation of triple-root jump in spacecraft charging, *J. Geophys. Res.*, Vol.96, No.A11, 19269-19282, 1991a

Lai, S.T., Spacecraft charging thresholds in single and double Maxwellian space environments, *IEEE Trans. Nucl. Sci.*, Vol.19, pp.1629-1634, 1991b

Laframboise J.G., Godard R., and Kamitsuma, Multiple floating potentials, "threshold-temperature" effects, and "barrier" effects in high-voltage charging of exposed surfaces in spacecraft, in *Proc Intl. Symp. Spacecraft Materials in Space Environment*, European Space Agency, ESA SP-178, 269-275,

1982.

Lauriente, M., A.L. Vampola, R. Koga and R. Hosken, Spacecraft anomalies due to the radiation environment, *AIAA 36th Aerospace Science Mtg.*, Reno, NV, Jan, 1998.

Robbins, A., Meteosat spacecraft charging investigation, European Space Agency, ESA CR(X) 1346, 1979.

Violet, M.D. and A.R. Frederickson, Spacecraft anomalies on the CRRES satellite correlated with the environment and insulator samples, *IEEE Trans. Nucl. Sci.*, Dec., 1993.

Whipple E.C., Jr., Potential of surface in space, *Rep. Prog. Phys.*, 44, 1197, 1981.

Wrenn, G.L., Conclusive evidence for internal dielectric charging anomalies on geosynchronous communications spacecraft. *J. Spacecraft & Rocket*, Vol.32, 514-520, 1995.

Wrenn, G.L. and A.J. Sims, Internal charging in the outer zone and operational anomalies, in *Radiation Belts: Models and Standards*, AGU Geophysical Monograph 97, AGU, Washington, DC, 1996.

Vampola, A.L., Thick dielectric charging on high-altitude spacecraft, *J. Electrostatics*, Vol.20, 21-30, 1987.